

UNIT 1: Physical Geography

A scenic view of a rocky coastline. In the foreground, a sandy beach curves along the water's edge. To the right, a large, craggy rock formation features a natural sea stack archway. The ocean is a deep blue, and the sky is clear. The overall scene is a beautiful natural landscape.

This exam is 1hr 30m long and is out of 88

It has 3 sections and 5 questions (you only answer Q1-4)

- Section A – Natural Hazards Q1 – Answer ALL questions (33/88)
- Section B – Living World Q2 – Answer ALL questions (25/88)
- Section C – Physical Landscapes Q3-5 – Answer Q3 & 4 ONLY
(each Q worth 15/88)

What are Natural Hazards?

Natural hazards are physical events such as earthquakes and volcanoes that have the potential to do damage to humans and property. Hazards include tectonic hazards, tropical storms and forest fires.


What affects hazard risk?

Population growth
Global climate change
Deforestation
Wealth - LICs are particularly at risk as they do not have the money to protect themselves



Structure of the Earth


The earth has 4 layers
The core (divided into inner and outer), mantle and crust.





The crust is split into major sections called **tectonic plates**.

Plates either move towards each other (**destructive margin**) away from each other (**constructive**) or past each other (**conservative**).

There are 2 types of crust: **Oceanic** (thin and younger but dense) and **Continental** (old and thicker but less dense).



These plates move due to convection currents in the mantle and, where they meet, tectonic activity (volcanoes and earthquakes) occurs..



Earthquakes and Volcanoes

Volcanoes

- **Constructive margins** – Hot magma rises between the plates e.g. Iceland. Forms Shield volcanoes.
- **Destructive margins** – an oceanic plate subducts under a continental plate. Friction causes oceanic plate to melt and pressure forces magma up to form composite volcanoes e.g. the west coast of South America.

Earthquakes

- **Constructive margins** – usually small earthquakes as plates pull apart.
- **Destructive margins** – violent earthquakes as pressure builds and is then released.
- **Conservative margins** – plates slide past each other. They catch and then as pressure builds it is released e.g. San Andreas fault.

Effects of Tectonic Hazards

Primary effects happen immediately. Secondary effects happen as a result of the primary effects and are therefore often later.

Primary - Earthquakes	Secondary - Earthquakes
<ul style="list-style-type: none"> - Property and buildings destroyed. - People injured or killed. - Ports, roads, railways damaged. - Pipes (water and gas) and electricity cables broken. 	<ul style="list-style-type: none"> - Business reduced as money spent repairing property. - Blocked transport hinders emergency services. - Broken gas pipes cause fire. - Broken water pipes lead to a lack of fresh water.

Primary - Volcanoes	Secondary - Volcanoes
<ul style="list-style-type: none"> - Property and farm land destroyed. - People and animals killed or injured. - Air travel halted due to volcanic ash. - Water supplies contaminated. 	<ul style="list-style-type: none"> - Economy slows down. Emergency services struggle to arrive. - Possible flooding if ice melts Tourism can increase as people come to watch. - Ash breaks down leading to fertile farm land.

Responses to Tectonic Hazards

Immediate (short term)	Long-term
<ul style="list-style-type: none"> - Issue warnings if possible. - Rescue teams search for survivors. - Treat injured. - Provide food and shelter, food and drink. - Recover bodies. - Extinguish fires. 	<ul style="list-style-type: none"> - Repair and re-build properties and infrastructure. - Improve building regulations - Restore utilities. - Resettle locals elsewhere. - Develop opportunities for recovery of economy. - Install monitoring technology.



Unit 1a The Challenge of Natural Hazards



Distribution of tectonic activity

Along plate boundaries.
On the edge of continents.
Around the edge of the Pacific.




Comparing Earthquakes – Haiti (LIC) and Kobe (HIC)

Haiti. Jan 2010. Magnitude 7.0.	Kobe Jan 1995. Magnitude 7.2.
Primary Effects	
Deaths 316,000 Injured 300,000 Wide scale devastation – presidential palace collapsed so little hope for those living in slums and shanty towns e.g. Cite Soleil Roads blocked by rubble Cost of damage \$30bn	Deaths 5,000 Newer buildings earthquake proof, but 102,000 older buildings collapsed Electricity and water supplies disrupted Phone communications disrupted Major expressway collapsed Cost of damage \$100bn
Secondary Effects	
Looting People forced into tented shelters Strong aftershocks Disease (Cholera) spread Damaged transportation	Fires from broken gas mains Homeless moved into well-built shelters The economy suffered as there was \$220 billion in damage. Companies like Panasonic had to close temporarily.
Immediate Responses	
Haiti needed foreign workers to help USAID with personnel, rescue dogs, and cutting equipment \$100m in aid given by USA and \$330m by EU UN flying in emergency food supplies Oxfam sending clean water, sanitation and shelter 4.3 million people provided with food rations	Government well prepared for earthquakes Japanese troops sent to help the people immediately Water, electricity, gas services were fully working by July 1995 Major retailers gave supplies to people affected Motorola maintained free mobile comms
Long term responses	
100m by World Bank to help with rebuilding 200,000 people received cash or food for clearing rubble	New buildings even more earthquake proof. More instruments to monitor earthquake movements

LICs suffer more than HICs from natural disasters because they are not as prepared and struggle to react effectively.

Global atmospheric circulation

At the equator, the sun's rays are most concentrated. This means it is hotter. This one fact causes global atmospheric circulation at different latitudes.



High pressure = dry
Low pressure = wet
As the air heats it rises – causing low pressure. As it cools, it sinks, causing high pressure. Winds move from high pressure to low pressure. They curve because of the **Coriolis effect** (the turning of the Earth)

Reducing the impact of tectonic hazards

Monitoring	Prediction
Seismometers measure earth movement. Volcanoes give off gases.	By observing monitoring data, this can allow evacuation before event.
Protection	Planning
Reinforced buildings and making building foundations that absorb movement. Automatic shut off for gas and electricity.	Avoid building in at risk areas. Training for emergency services and planned evacuation routes and drills.


Tropical Storms

Occur in low latitudes between 5° and 30° north and south of the equator (in the tropics). Ocean temperature needs to be above 27° C. Happen between summer and autumn.



Sequence of a Tropical Storm

- Air is heated above warm tropical oceans.
- Air rises under low pressure conditions.
- Strong winds form as rising air draws in more air and moisture causing torrential rain.
- Air spins due to Coriolis effect around a calm eye of the storm.
- Cold air sinks in the eye so it is clear and dry.
- Heat is given off as it cools powering the storm.
- On meeting land, it loses source of heat and moisture so loses power.



Climate change will affect tropical storms too. Warmer oceans will lead to more intense storms – but not necessarily more frequent ones.

Extreme weather in the UK

- Rain** – can cause flooding damaging homes and business.
- Snow & Ice** – causes injuries and disruption to schools and business. Destroys farm crops.
- Hail** – causes damage to property and crops.
- Drought** – limited water supply can damage crops.
- Wind** – damage to property and damage to trees potentially leading to injury.
- Thunderstorms** – lightning can cause fires or even death.
- Heat waves** – causes breathing difficulties and can disrupt travel.

UK weather is getting more extreme due to climate change. Temperatures are more extreme and rain is more frequent and intense leading to more flooding events. Since 1980 average temperature has increased 1 degree and winter rainfall has increased.

Hurricane Katrina, USA, August 2005



Primary Effects	Secondary Effects
At least 1800 killed 120 mph wind speeds 300,000 houses destroyed Levees broke Coastal habitats damaged 80% New Orleans flooded	\$150bn of damage Water supply polluted 230,000 jobs lost from damaged businesses 200,000 people made homeless Dehydration of people awaiting rescue

Immediate Responses	Long-term Responses
70-80% of New Orleans evacuated before the hurricane struck Mississippi & Louisiana declared states of emergency and set up control centres, emergency shelters & supplies Coastguard, police, fire and army rescued 50,000 people Charities gave aid including millions of hot meals	US Government gave \$16bn for rebuilding homes and funds for other infrastructure New homes built on stilts or not at all in high risk areas Repaired and improved flood defences costing \$14.5bn

Prediction	Planning	Protection
Monitoring wind patterns allows path to be predicted. Use of satellites to monitor path to allow evacuation	Avoid building in high risk areas Emergency drills Evacuation routes	Reinforced buildings and stilts to make safe Flood defences e.g. levees and sea walls Replanting Mangroves

Feb 2018 – The Beast from the East

A long period of heavy snow and cold weather across the UK because of cold air from northern Europe and Siberia

Social Effects

10 people died from hypothermia or accidents on icy roads
Shortage of food in supermarkets
Thousands of schools closed on several occasions meaning parents take time off work

Economic Effects

Overall economic impact was approx. £1bn a day
Transport (roads/rail/air) networks closed causing businesses to lose money
People unable to get to work
Construction industry the biggest hit

Environmental impacts

Snow was up to 50cm deep with gusts of wind up to 70mph
The frost killed crops
Amount of gas & electricity used to heat homes went up from normal use increasing CO₂ emissions

Management strategies

Met Office issued a red weather warning of cold weather in Feb
Councils stocked up on gritters and salt supplies to keep roads open and safe (though not enough)
Military personnel drafted in to rescue people stranded in cars and delivered medical personnel

Climate Change – natural or human?

Evidence for climate change shows changes before humans were on the planet. So some of it must be natural. However, the **rate** of change since the 1970s is unprecedented. Humans are responsible – despite what Mr Trump says!

Causes


Natural	Human
<ul style="list-style-type: none"> Orbital changes – The sun's energy on the Earth's surface changes as the Earth's orbit is elliptical its axis is tilted on an angle. Solar Output – sunspots increase to a maximum every 11 years. Volcanic activity – volcanic aerosols reflect sunlight away reducing global temperatures temporarily. 	<ul style="list-style-type: none"> Fossil fuels – release carbon dioxide with accounts for 50% of greenhouse gases. Agriculture – accounts for around 20% of greenhouse gases due to methane production from cows etc. Larger populations and growing demand for met and rice increase contribution. Deforestation – logging and clearing land for agriculture increases carbon dioxide in the atmosphere and reduces ability to planet to absorb carbon through photosynthesis.

Effects of Climate Change

Social	Environmental
<ul style="list-style-type: none"> Increased disease e.g. skin cancer and heat stroke. Winter deaths decrease with milder winters. Crop yields affected by up to 12% in South America but will increase in Northern Europe but will need more irrigation. Less ice in Arctic Ocean increases shipping and extraction of oil and gas reserves. Droughts reduce food and water supply in sub-Saharan Africa. Water scarcity in South and South East UK. Increased flood risk. 70% of Asia is at risk of increased flooding Declining fish in some areas affect diet and jobs. Increased extreme weather Skiing industry in Alps threatened. 	<ul style="list-style-type: none"> Increased drought in Mediterranean region. Lower rainfall causes food shortages for orangutans in Borneo and Indonesia. Sea level rise leads to flooding and coastal erosion. Ice melts threaten habitats of polar bears. Warmer rivers affect marine wildlife. Forests in North America may experience more pests, disease and forest fires. Coral bleaching and decline in biodiversity.

Managing Climate Change

Mitigation	Adaptation
<ul style="list-style-type: none"> Alternative energy production will reduce CO₂ production. Planting Trees – helps to remove carbon dioxide. Carbon Capture – takes carbon dioxide from emission sources is stored underground. International Agreements e.g. the Paris Climate Agreement. 	<ul style="list-style-type: none"> Changes in agricultural systems need to react to changing rainfall and temperature patterns and threat of disease and pests. Managing water supplies – e.g. by installing water efficient devices and increasing supply through desalination plants. Reducing risk from rising sea levels would involve constructing defences such as the Thames Flood Barrier or restoring mangrove forests, or raising buildings on stilts.



Evidence for Climate Change

The Met Office has reliable climate evidence since 1914 – but we can tell what happened before that using several methods.

Ice and Sediment Cores

- Ice sheets are made up of layers of snow, one per year. Gases trapped in layers of ice can be analysed. Ice cores from Antarctica show changes over the last 400 000 years.
- Remains of organisms found in cores from the ocean floor can be traced back 5 million years.

Pollen Analysis

Pollen is preserved in sediment. Different species need different climatic conditions.

Tree Rings

- A tree grows one new ring each year. Rings are thicker in warm, wet conditions
- This gives us reliable evidence for the last 10 000 years.

Temperature Records

Historical records date back to the 1850s. Historical records also tell us about harvest and weather reports.

